


Factors influencing carbon storage capacity in eelgrass meadows of New England



**Alyssa Novak¹, Phil Colarusso², Marguerite Pelletier³,
Julie Simpson⁴, Mary Nicole Gutierrez⁵, Ariane Arias
Ortiz⁶, Pere Masqué⁶, Prasseda Vella⁷**

¹Boston University, Earth and Environment, Boston, MA, USA, ²US EPA, Region 1, Boston, MA, USA, ³US EPA, ORD, NHEERL-AED, Narragansett, RI, USA, USA,

⁴ MIT Sea Grant, Cambridge, MA, USA, ⁵ SSC to US EPA, Narragansett, RI, USA,

⁶ Autonomous University of Barcelona, Barcelona, Spain, ⁷ Mass CZM, Boston, MA, USA

Multimodel Inference

- Alternative to null hypothesis testing
- Multiple models generated using different combinations of variables
- Information criteria (AIC) used to rank models
- Relative importance of each variable in model is determined from AIC_w value summed across models

Multimodel Inference

- Alternative to null hypothesis testing
- Multiple models generated using different combinations of variables
- Information criteria (AIC) used to rank models
- Relative importance of each variable in model is determined from AIC_w value summed across models

Multimodel Inference Conceptual model

Variables

- Environmental Conditions
 - Latitude (temperature)
 - Fetch
 - Precipitation
- Sediment Characteristics
 - % Silt Clay
 - Percent %N
 - Trap weight
- Plant Parameters
 - Leaf Area
 - Growth Rate (cm)
 - Shoot Density



Multimodel Inference Conceptual model

Variables

- Environmental Conditions
 - Latitude (temperature)
 - Fetch
 - Precipitation
- Sediment Characteristics
 - % Silt Clay
 - Percent %N
 - Trap weight
- Plant Parameters
 - Leaf Area
 - Growth Rate (cm)
 - Shoot Density



Multimodel Inference Conceptual model

Variables

- Environmental Conditions
 - Latitude (temperature)
 - Fetch
 - Precipitation
- Sediment Characteristics
 - % Silt Clay
 - Percent %N
 - Trap weight
- Plant Parameters
 - Leaf Area
 - Growth Rate (cm)
 - Shoot Density



Multimodel Inference Conceptual model

Variables

- Environmental Conditions
 - Latitude (temperature)
 - Fetch
 - Precipitation
- Sediment Characteristics
 - % Silt Clay
 - Percent %N
 - Trap weight
- Plant Parameters
 - Leaf Area
 - Growth Rate (cm)
 - Shoot Density




Multimodel Inference Conceptual model

Top 5 cm- Eelgrass

- % silt-clay= high importance
- Latitude & %N trap = low to moderate importance
- Remaining variables= low importance

Model has moderate predictive ability

Variable	Sum AICw
% silt-clay	0.86
Latitude	0.29
Sed trap %N	0.27
Leaf area	0.18
Avg growth	0.14
Total spring/summer ppt	0.13
Avg shoot density	0.11
sed trap wt	0.11
Median fetch	0.10




Multimodel Inference Conceptual model

Top 5 cm- Eelgrass

- % silt-clay= high importance
- Latitude & %N trap = low to moderate importance
- Remaining variables= low importance

Model has moderate predictive ability

Variable	Sum AICw
% silt-clay	0.86
Latitude	0.29
Sed trap %N	0.27
Leaf area	0.18
Avg growth	0.14
Total spring/summer ppt	0.13
Avg shoot density	0.11
sed trap wt	0.11
Median fetch	0.10




Multimodel Inference Conceptual model

Top 5 cm- Eelgrass




- % silt-clay= high importance
- Latitude & %N trap = low to moderate importance
- Remaining variables= low importance

Model has moderate predictive ability

Variable	Sum AICw
% silt-clay	0.86
Latitude	0.29
Sed trap %N	0.27
Leaf area	0.18
Avg growth	0.14
Total spring/summer ppt	0.13
Avg shoot density	0.11
sed trap wt	0.11
Median fetch	0.10






Multimodel Inference Conceptual model



Top 5 cm- Eelgrass

Multimodel Inference Conceptual model



Top 5 cm- Eelgrass


Multimodel Inference Conceptual model

Top 5 cm- Reference

- % silt-clay= high importance
- Latitude & Precipitation= low to moderate importance
- Remaining variables= low importance

Model has good predictive ability

Variable	Sum AICw
% silt-clay	0.96
Latitude	0.25
Total spring/summer ppt	0.23
Median fetch	0.16
sed trap wt	0.16
Sed trap %N	0.14




Multimodel Inference Conceptual model

Top 5 cm- Reference




- % silt-clay= high importance
- Latitude & Precipitation= low to moderate importance
- Remaining variables= low importance

Model has good predictive ability

Variable	Sum AICw
% silt-clay	0.96
Latitude	0.25
Total spring/summer ppt	0.23
Median fetch	0.16
sed trap wt	0.16
Sed trap %N	0.14




Multimodel Inference Conceptual model



Top 5 cm- Reference


Future: Relative Wave Exposure

- WEMo- NCCOS: uses linear wave theory to calculate wave height and wave energy
 - wind generation
 - water depth




Future: Relative Wave Exposure

- WEMo- NCCOS: uses linear wave theory to calculate wave height and wave energy
 - wind generation
 - water depth



Future: Relative Wave Exposure

- WEMo- NCCOS: uses linear wave theory to calculate wave height and wave energy
 - wind generation
 - water depth



Conclusions

- We have developed reasonable models with moderate-good predictive ability with collected variables.
- Sediment grain size is an important predictor while latitude is a moderate predictor of bulk carbon at eelgrass and reference sites.
- Future models will substitute fetch with relative wave exposure.

Acknowledgements

Fred Short, Eric Nelson, Dan Arsenault, Jean Brochi, Chuck Protzmann, Tim Bridges, Holly Plaisted, Agnes Mittermayr, Jill Carr, Tay Evans, Jack Buckley, Dani Ewart, Dave Grunden, Hanna Mogensen, Juliette Williams, Lindsay Peter, Audrey Michniak, Molly Sullivan, Ivy Mlsna, Kathryn Baltes, Bill Osbahr, Don Cobb

• .

